

Chapter 5

Applying Newton's Laws

PowerPoint® Lectures for
University Physics, Thirteenth Edition
– Hugh D. Young and Roger A. Freedman

Lectures by Wayne Anderson

Copyright © 2012 Pearson Education Inc.

Goals for Chapter 5

- To use Newton's first law for bodies in equilibrium
- To use Newton's second law for accelerating bodies
- To study the types of friction and fluid resistance
- To solve problems involving circular motion

Copyright © 2012 Pearson Education Inc.

Introduction

- We'll start with equilibrium, in which a body is at rest or moving with constant velocity.
- Next, we'll study objects that are not in equilibrium and deal with the relationship between forces and motion.
- We'll analyze the friction force that acts when a body slides over a surface.
- We'll analyze the forces on a body in circular motion at constant speed.

Copyright © 2012 Pearson Education Inc.

Particles in Equilibrium

If the acceleration of an object that can be modeled as a particle is zero, the object is said to be in **equilibrium**

- The model is the *particle in equilibrium model*

Mathematically, the net force acting on the object is zero

$$\sum \vec{F} = 0$$
$$\sum F_x = 0 \text{ and } \sum F_y = 0$$

Copyright © 2012 Pearson Education Inc.


Two-dimensional equilibrium : Example

- A car engine hangs from several chains.
- Find the expression for the tension in each of three chains in terms of w

(a) Engine, chains, and ring

(b) Free-body diagram for engine


(c) Free-body diagram for ring O



Copyright © 2012 Pearson Education Inc.

Equilibrium, Example 2

Find tension in each cable if the weight of street lamp is 122 N



© 2012 Pearson Higher Education

Copyright © 2012 Pearson Education Inc.


Inclined Planes

Forces acting on the object:

- The normal force acts perpendicular to the plane
- The gravitational force acts straight down

Choose the coordinate system with x along the incline and y perpendicular to the incline

Replace the force of gravity with its components



Copyright © 2012 Pearson Education Inc.

A car on an inclined plane


An car of weight w rests on a slanted ramp attached to a trailer. Only a cable running from the trailer prevents the car from rolling off the ramp. Find the tension in the cable and the force that the ramp exerts on the car's tires. $W = mg$

$$T - mgsina = 0 \quad T = mgsina$$

$$n - mgcosa = 0 \quad n = mgcosa$$

(a) Car on ramp


(b) Free-body diagram for car



Copyright © 2012 Pearson Education Inc.

Incline plane. Object not in equilibrium: $\Sigma F = ma$

- What is the acceleration of a toboggan loaded with students of total mass of 500 kg sliding down a friction-free slope? Assume $\alpha = 15^\circ$.




Copyright © 2012 Pearson Education Inc.

Multiple Objects. Bodies connected by a cable and pulley

A 500-N cart is connected to a bucket by a cable passing over a pulley. What mass of the bucket is needed for the system to move up the 15° incline with constant speed?

- Draw separate free-body diagrams for the bucket and the cart.
- Apply Newton's second law $\Sigma F_x = ma_x$, $\Sigma F_y = ma_y$ for each element of the system.




Copyright © 2012 Pearson Education Inc.

Multiple Objects, Example

Find the magnitude of the acceleration of the system if $m_1 = 2\text{ kg}$, $m_2 = 5.5\text{ kg}$, incline angle $\theta = 32^\circ$.

Draw the free-body diagram for each object


- One cord, so tension is the same for both objects
- Connected, so acceleration is the same for both objects




Copyright © 2012 Pearson Education Inc.

Frictional forces

- When a body rests or slides on a surface, the friction force is parallel to the surface.



- Friction between two surfaces arises from interactions between molecules on the surfaces.



On a microscopic level, even smooth surfaces are rough; they tend to catch and cling.

Copyright © 2012 Pearson Education Inc.


Kinetic and static friction

- Kinetic friction acts when a body slides over a surface.
- The kinetic friction force is $f_k = \mu_k n$.
- Static friction acts when there is no relative motion between bodies.
- The static friction force can vary between zero and its maximum value: $f_s \leq \mu_s n$.

Copyright © 2012 Pearson Education Inc.

Static friction followed by kinetic friction

- Before the box slides, static friction acts. But once it starts to slide, kinetic friction acts.



Copyright © 2012 Pearson Education Inc.

Some approximate coefficients of friction

Table 5.1 Approximate Coefficients of Friction


Materials	Coefficient of Static Friction, μ_s	Coefficient of Kinetic Friction, μ_k
Steel on steel	0.74	0.57
Aluminum on steel	0.61	0.47
Copper on steel	0.53	0.36
Brass on steel	0.51	0.44
Zinc on cast iron	0.85	0.21
Copper on cast iron	1.05	0.29
Glass on glass	0.94	0.40
Copper on glass	0.68	0.53
Teflon on Teflon	0.04	0.04
Teflon on steel	0.04	0.04
Rubber on concrete (dry)	1.0	0.8
Rubber on concrete (wet)	0.30	0.25

Copyright © 2012 Pearson Education Inc.

Friction in horizontal motion

To start a 500-N crate moving across a level floor you have to pull with a 230-N horizontal force. Once the crate starts to move, you can keep it moving at constant speed with only 200N. What are the coefficients of static and kinetic friction?


Before the crate moves, static friction acts on it. After it starts to move, kinetic friction acts.



Copyright © 2012 Pearson Education Inc.

Friction - Example


Find the coefficient of static friction force if the block starts to slide down at an angle of 12.5°



Copyright © 2012 Pearson Education Inc.

Motion on a slope having friction


What is the acceleration of a toboggan loaded with students of total mass of 500 kg sliding down a slope? Assume $\alpha = 15^\circ$ and coefficient of kinetic friction the slope $\mu_k = 0.08$.



Copyright © 2012 Pearson Education Inc.


Friction, Example

A hockey puck on a frozen pond is given an initial speed of 12 m/s. If the puck slides 60 m before coming to rest, determine the coefficient of kinetic friction between the puck and ice.



Copyright © 2012 Pearson Education Inc.

Friction, Example




Find the acceleration of the system of $m_1 = 5 \text{ kg}$, $m_2 = 1.5 \text{ kg}$, $F = 25 \text{ N}$, $\mu_k = 0.12$, and $\theta = 28^\circ$

Copyright © 2012 Pearson Education Inc.

Pulling a crate at an angle

- The angle of the pull affects the normal force, which in turn affects the friction force.
- How hard must you pull the crate to keep it moving with constant velocity? Assume $\mu_k = 0.4$





Copyright © 2012 Pearson Education Inc.

Fluid resistance and terminal speed

- The *fluid resistance* on a body depends on the speed of the body.
- A falling body reaches its *terminal speed* when the resisting force equals the weight of the body.
- The figures at the right illustrate the effects of air drag.


(a) Free-body diagrams for falling with air drag



Copyright © 2012 Pearson Education Inc.

Dynamics of circular motion

- If a particle is in uniform circular motion, both its acceleration and the net force on it are directed toward the center of the circle.
- The net force on the particle is $F_{\text{net}} = mv^2/R$.




Copyright © 2012 Pearson Education Inc.

What if the string breaks?

- If the string breaks, no net force acts on the ball, so it obeys Newton's first law and moves in a straight line.

A ball attached to a string whirls in a circle on a frictionless surface.




No net force now acts on the ball, so it obeys Newton's first law—it moves in a straight line at constant velocity.

Copyright © 2012 Pearson Education Inc.


Force in uniform circular motion

- A sled on frictionless ice is kept in uniform circular motion by a rope.
- $F =$

(a) A sled in uniform circular motion



(b) Free-body diagram for sled




Copyright © 2012 Pearson Education Inc.


A conical pendulum

A bob at the end of a wire moves in a horizontal circle with constant speed.

(a) The situation



(b) Free-body diagram for pendulum bob




Copyright © 2012 Pearson Education Inc.


A car rounds a flat curve

- A car rounds a flat unbanked curve of radius $R=65$ m. What is its maximum speed if the $\mu_s=0.58$?

(a) Car rounding flat curve



(b) Free-body diagram for car




Copyright © 2012 Pearson Education Inc.


A car rounds a banked curve

- At what angle should a curve be banked so a car can make the turn even with no friction?

(a) Car rounding banked curve



(b) Free-body diagram for car



Copyright © 2012 Pearson Education Inc.


Uniform motion in a vertical circle

- A person on a Ferris wheel moves in a vertical circle.


Top: $n-w = -ma_c$ $n = w - ma_c$

Bottom: $n-w = ma_c$ $n = w + ma_c$


(a) Sketch of two positions



(b) Free-body diagram for passenger at top



(c) Free-body diagram for passenger at bottom



Copyright © 2012 Pearson Education Inc.